



BIRDS



Wild fowl maintenance

Feed number: 3972

Suitable for: Park, wild and ornamental fowl

YOUR BENEFITS

- ◇ Balanced complete feed for the supply of all nutrients, minerals, trace elements and vitamins
- ◇ Fish oil for natural omega 3 fatty acids
- ◇ Convenient granules for an easy feed intake
- ◇ With phytase to improve phosphorus utilization (6-phytase: 750 FTU/kg)



Full-scale picture of the feed
Colours may be different from the product

TYPE OF FEED, FORM, DELIVERY QUANTITY

- ◇ Complete feed
- ◇ Form: granules
- ◇ Delivery quantity: 25 kg paper bag
pallets of 750 kg
- ◇ Product number: 3972.CI.S25



Wild fowl maintenance

Feed number: 3972

FEED SPECIFICATIONS

Major nutrients (%)

Dry matter	88
Crude protein	15
Crude fat	3
Crude fiber	7
Crude ash	7
NFE	56
NDF	18
ADF	8
Starch	41
Sugar	3

Energy (MJ/kg)

GE	16
ME	10.6

Macrominerals (%)

Calcium	1.2
Phosphorus	0.6
Magnesium	0.2
Sodium	0.2
Potassium	0.8
Chlorine	0.2

Trace elements (mg/kg)

Iron	350
Zinc	96
Copper	17
Iodine	1.7
Manganese	120
Selenium	0.4
Cobalt	0.1

Vitamins (added, mg/kg)

Vitamin A (IU/KG)	9500
Vitamin D3 (IU/KG)	3000
Vitamin E	40
Vitamin K3	4
Vitamin B1	3
Vitamin B2	12
Vitamin B6	5
Vitamin B12	0.03
Nicotinic acid	70
Pantothenic acid	30
Folic acid	1.7
Biotin	0.3
Choline	400
Vitamin C	30

Amino acids (%)

Arginine	0.88
Lysine	0.03
Methionine	0.48
Methionine + Cystine	0.80
Tryptophan	0.18
Threonine	0.52

Ingredients

Wheat, corn (NGMO), oats, wheat bran, lucerne, soybean meal (NGVO), protein peas, calcium carbonate, corn gluten (NGMO), mineral and vitamin premixes, soybean oil (NGMO), fish oil, amino acids

Remarks

- ◇ Given values are calculated averages in air-dry feed.
- ◇ Energy values calculated according to Kamphues et al. 2014. GE=gross energy, ME=metabolisable energy.
- ◇ Trace elements: calculated total content. Vitamin declaration: vitamins added before production. Estimated total vitamin content for ration calculations on request.
- ◇ Nutrients are subject to natural variation of the raw materials and their production process.

OUR FEED RECOMMENDATION

- ◇ This feed is offered ad libitum to the animals.
- ◇ Suitable for birds from the 8th-10th week of life
- ◇ Add the Wildfowl breeding 4-6 weeks before the onset of egg laying
- ◇ Always provide free access to fresh drinking water.

Sources:

Kamphues, J. et al. (2014). Supplemente zu Vorlesungen und Übungen in der Tierernährung. 18. National Research Council (1994), Nutrient Requirements of poultry, Ninth ed.